

Advice to Householders



Your property has been treated with rodenticide today by:
Section to be completed by operator completing treatment.

Operators Name: _____	Product Used: _____
Operators Company: _____	Date of Treatment: _____
Company Phone Number: _____	

Contains The Following Active Ingredient: (Operator to tick appropriate box)

FLOCOUMAFEN

DIFENACOUM

THIS PRODUCT CONTAINS AN ANTICOAGULANT.

You are advised that this rodenticide is approved for use in domestic premises, e.g. houses and flats, under the Control of Pesticides Regulations 1986. However, your co-operation is requested in adopting the following precautions to further minimise any risks to occupants and pets.

Precautions

AVOID ALL CONTACT BY MOUTH.

PREVENT ACCESS TO BAIT

by children, birds and non-target animals (particularly dogs, cats, pigs and poultry).

IN CASE OF ACCIDENT,

contact a doctor or veterinarian and show this leaflet.

**DO NOT
MOVE OR
TAMPER
WITH BAIT.**

Advice to Physicians

If you think that bait has been eaten – The product is an indirect anticoagulant. Phytomenadione, Vitamin K1, is antidotal. Determine prothrombin times not less than eighteen hours after consumption. If elevated, administer Vitamin K1 in divided doses. Continue until prothrombin times normalise. After withdrawal of the antidote, continue determination of prothrombin time for two weeks and resume treatment if elevation occurs in that time. NB. Vitamin K3 is not effective.

For comprehensive medical advice on the treatment of poisoning, contact the nearest Poisons Information Centre or BASF.

Please note: BASF has not carried out this treatment. This form has been provided by BASF to the operator making your treatment. Please contact them, regarding any questions on your treatment.

**BASF Plc
Pest Control Solutions**

PO Box 4, Earl Road, Cheadle Hulme, Cheadle, Cheshire, SK8 6QG, United Kingdom
e-mail: pestinfo@basf.com www.pestcontrol.basf.co.uk

Declaration: BASF accepts no liability if any amendments are made to this leaflet.

Advice to Householders



Your property has been treated with rodenticide today by:
Section to be completed by operator completing treatment.

Operators Name: _____	Product Used: _____
Operators Company: _____	Date of Treatment: _____
Company Phone Number: _____	

Contains The Following Active Ingredient: (Operator to tick appropriate box)

FLOCOUMAFEN

DIFENACOUM

THIS PRODUCT CONTAINS AN ANTICOAGULANT.

You are advised that this rodenticide is approved for use in domestic premises, e.g. houses and flats, under the Control of Pesticides Regulations 1986. However, your co-operation is requested in adopting the following precautions to further minimise any risks to occupants and pets.

Precautions

AVOID ALL CONTACT BY MOUTH.

PREVENT ACCESS TO BAIT

by children, birds and non-target animals (particularly dogs, cats, pigs and poultry).

IN CASE OF ACCIDENT,

contact a doctor or veterinarian and show this leaflet.

**DO NOT
MOVE OR
TAMPER
WITH BAIT.**

Advice to Physicians

If you think that bait has been eaten – The product is an indirect anticoagulant. Phytomenadione, Vitamin K1, is antidotal. Determine prothrombin times not less than eighteen hours after consumption. If elevated, administer Vitamin K1 in divided doses. Continue until prothrombin times normalise. After withdrawal of the antidote, continue determination of prothrombin time for two weeks and resume treatment if elevation occurs in that time. NB. Vitamin K3 is not effective.

For comprehensive medical advice on the treatment of poisoning, contact the nearest Poisons Information Centre or BASF.

Please note: BASF has not carried out this treatment. This form has been provided by BASF to the operator making your treatment. Please contact them, regarding any questions on your treatment.

**BASF Plc
Pest Control Solutions**

PO Box 4, Earl Road, Cheadle Hulme, Cheadle, Cheshire, SK8 6QG, United Kingdom
e-mail: pestinfo@basf.com www.pestcontrol.basf.co.uk

Declaration: BASF accepts no liability if any amendments are made to this leaflet.