



RODILON®

Wax Blocks

Rodenticide

RODILON® RAT AND MOUSE WAX BLOCKS

Rodenticide - Single-Feed

Reg. No. L5356 Act No. 36 of 1947

Before using this product read the label carefully.

Active anticoagulant in block form for control of the Norway rat, roof rat and house mouse. Effective against warfarin resistant rats and mice. For use in the home, on the farm, in public health and industrial premises.

Hazard statements
May cause damage to organs (Blood) through prolonged or repeated exposure. To avoid risks to human health and the environment, comply with the instructions for use.
Precautionary statements
Do not breathe dust.
Avoid release to the environment.
Wear protective gloves.



Warning

ACTIVE INGREDIENT:
Difethialone (anticoagulant)0,025 g/kg

REGISTRATION HOLDER:
2022 Environmental Science ZA (Pty) Ltd. Reg. No. 2022/228577/07
P.O. Box 143, ISANDO 1600, SOUTH AFRICA
TEL: (011) 921 5911

RODILON® is a registered trademark of the Bayer Group
* A Lipha Product

10 kg



Norway Rat . House Mouse

D00000420A

DISCLAIMER:

ALTHOUGH THIS PRODUCT HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS, FAILURE THEREOF MAY ENSUE AS A RESULT OF A WIDE RANGE OF REASONS. IF THIS IS SUSPECTED, KINDLY NOTIFY THE REGISTRATION HOLDER.

WARNINGS:

Hazard statements: Harmful to aquatic life with long lasting effects.
Handle with care. Keep out of reach of children, uninformed persons and animals. Poisonous if swallowed and upon prolonged contact with the skin. Toxic to fish and wildlife. Store away from food and feedstuffs.
Do not place the bait at random in treated area but use exclusively as given under directions for use. In case of poisoning, call a doctor and make the insert available to them. Shelf-Life: expires 2 years after date of manufacture.

PRECAUTIONS:

Precautionary statements: Keep out of reach of children. Do not eat, drink or smoke when using this product. Get medical advice/ attention if you feel unwell.
Dispose of contents/container in accordance with local regulation. Avoid contact with skin. Wear gloves when handling product. Avoid contamination of any food, food utensils and drinking water. Set bait where it will be inaccessible to children and animals. Wash hands and exposed skin after handling. The use of bait boxes or other special containers is strongly recommended. Dispose of dead rats, any bait not consumed and empty containers in accordance with current regulations and, if necessary, after consultation with the site operator and/or with the responsible authority, the product may be taken to a waste disposal site or incineration plant.
RELEVANT SUBSTANCES:

Name	CAS-No. / EC-No. / REACH Reg. No.	Conc. [%]
difethialone	104653-34-1	0,0025

SYMPTOMS OF POISONING:

Bleeding from the gums and nose, bleeding under the skin, pallor, haemorrhage into and around elbow and knee joints. Blood in urine and faeces.

FIRST AID MEASURES

General advice:	Move out of dangerous area. Place and transport victim in stable position (lying sideways). Remove contaminated clothing immediately and dispose of safely.
Skin contact:	Wash off thoroughly with plenty of water and soap, if available with polyethyleneglycol 400, subsequently rinse with water
Eye contact:	Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Get medical attention if irritation develops and persists.
Ingestion:	IMMEDIATELY remove to fresh air. If not breathing give artificial respiration. If breathing of victim is difficult administer oxygen for a maximum period of one hour. Have victim lie down and keep warm.

Have the product container or label with you when calling a poison control centre / doctor / going for treatment.

NOTE TO PHYSICIAN

Treatment Elementary aid, decontamination and symptomatic treatment.

Antidote Administer 5 mg/kg Vit. K, intravenously on the first day, followed by another administration of the same dose if the Quick Time does not normalise a few hours after the first injection. Follow up with 5 mg/kg/day orally on the second and following days. The duration of the treatment is generally between 10 and 15 days. In severe cases give small transfusions of fresh blood.

PRIMARY ANIMAL POISONING: In case of primary poisoning in animals, contact the veterinarian immediately.

NOTE TO VETERINARIAN:

SPECIFIC ANTIDOTE:

Vit K, is the only antidote in all cases of Coumarin anti-coagulant poisons.

ANIMALS EXHIBITING SIGNS OF INTOXICATION

Collect a blood sample and carry out a prothrombin test, Parenteral injection of Vitamin K, 2 to 5mg/kg. Use the smallest diameter needle feasible, and avoid the intravenous route in severely haemorrhagic animals. Supportive measures. Repeat prothrombin test about four hours after injection. Provided prothrombin time has normalised, start daily oral Vitamin K1 treatment and continue it for three to four weeks. Carry out prothrombin test 24 to 48 hours after end of treatment. Continue treatment if signs of poisoning reappear or if prothrombin time is still abnormal.

ANIMALS SUSPECTED OF CONSUMING BAIT

This includes animals known to have ingested the bait or animals showing evidence of ingestion such as traces of dye from the bait around the mouth or in faeces. Induce vomiting only if ingestion is recent (less than six hours). Observe closely for signs of poisoning for one week. Test prothrombin time daily for up to three days after suspected date of ingestion. Treat with Vitamin K1 if signs of poisoning appear or if prothrombin time increases.

Prophylactic oral Vitamin K1 treatment could be carried out.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

Rodilon® Rat and Wax Blocks can kill rats and mice after a single feed. Death of rodents will occur in 4 - 12 days.
Effective against warfarin resistant rats and mice.
For use on the farm and industrial premises (outside buildings, warehouses and stores).
Before laying bait inspect infested area thoroughly to determine rat or mouse entry, nesting and feeding sites and water sources

PESTS	DOSAGE RECOMMENDATIONS	REMARKS
Norway Rats and Roof Rats	Initial baiting 3-4 wax blocks per bait station (5-10 m apart)	Place Rodilon® Wax blocks in a covered bait station to prevent access by children and domestic animals.
	Maintenance baiting Place 1 block per bait station every 5 - 10 m (or 50 m ²) to maintain control	Place Rodilon® Wax blocks along walls or rat-runs in bait stations, in rat burrows and in sheltered areas where rats are known to live or feed. Bait points should be inspected and replenished every 7 days until feeding ceases.
House Mice	1 wax block per bait station (2-4 m apart)	Place Rodilon® Wax blocks per bait station along walls or runways in bait stations, in nests and in sheltered areas where mice are known to live or feed. The single feed action of Rodilon® Wax blocks is particularly effective against mice with their small appetites and sporadic feeding behaviour. If necessary, replenish every 7 - 14 days until feeding ceases. To prevent re-infestation, maintain a few permanent baiting points.

For any product enquiry, contact 2022 Environmental Science ZA (Pty) Ltd. - 011 921 5911 (Office Hours 9 am - 5pm)

For poisoning emergencies only, contact Griffon Poison Information Centre 082 446 8946.

RODILON® is a registered trademark of the Bayer Group.
© 2022 Environmental Science ZA (Pty) Ltd. All unauthorized reproduction of this label is forbidden.

Batch no: / Lotnummer:
Date of manufacture: / Datum vervaardig:
Shelf-Life: expires 2 years from date of manufacture.




Code: D00000420A Plan: XXX Size: 285 x 145 mm Date: 24/07/2023 Op: Pierwood/AL

C

M

Y

K